



Ethnobotanical uses and phytochemical, biological, and toxicological profiles of *Datura metel* L.: A review

Tawhida Islam^a, Iffat Ara^a, Tariqu Islam^a, Pankaj Kumar Sah^a, Ray Silva de Almeida^b,
Edinardo Fagner Ferreira Matias^c, Cícero Lucas Gomes Ramalho^c, Henrique Douglas
M. Coutinho^{b,*}, Muhammad Torequ Islam^{a,*}

^a Department of Pharmacy, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj 8100, Bangladesh

^b Department of Biological Chemistry, Regional University of Cariri—URCA, Crato 63105-000, Brazil

^c CECAPE - College of Dentistry, Juazeiro do Norte, CE 63024-015, Brazil

ARTICLE INFO

Keywords:

Datura metel
Poisonous plant
Phytochemicals
Biological activities

ABSTRACT

Datura metel L., a recognized poisonous plant in the Solanaceae family, is widely distributed in the world. Traditionally, *D. metel* is used in many diseases, including neurological and heart diseases; fever; catarrh; pain; diarrhea; skin diseases; chronic bronchitis; asthma; digestive disorders; and so on. It possesses many important phytochemicals that can be used to treat various types of diseases. This review aims at summarizing the traditional uses, phytochemical, biological, and toxicological profiles of *D. metel* based on the database reports. For this, an up-to-date (till March 20, 2023) search was made in the databases: PubMed, Google Scholar, Science Direct, Scopus, and MedLine, with relevant keywords for the published evidence. Findings suggest that the plant has many traditional uses, such as a cure for madness, epilepsy, psoriasis, heart diseases, diarrhea, mad dog bites, indigestion, etc. It possesses various important phytochemicals, including withanolides, daturaolone, datumetine, daturglycosides, ophiobolin A, baimantuoluoline A, and many others. *D. metel* has many important biological activities, including antioxidant, anti-inflammatory, anti-microbial, insecticidal, anti-cancer, anti-diabetic, analgesic, anti-pyretic, neurological, contraceptive, and wound healing capacity. In conclusion, the toxic plant, *D. metel*, can be considered a potential source of phyto-therapeutic lead compounds.

Introduction

In Bengali, the local name of *Datura* (*Datura metel* L.) is “Dhutura-ধুতুরা”. Other common names are in Arabic: tatura, jozmashel, jozmathel; in Chinese: yang jinhua; in English: purple thorn-apple, downy thorn-apple, Hindu datura, hoary thorn-apple, Hindu thorn-apple, horn-of-plenty; in Hindi: sadadhatura; in Korean: huindogmalpul; in Portuguese: burbiaca; in Spanish: burladora; in Sweden: indisk spikklubba (Al-Snafi, 2017a). The genus *Datura* (Family: Solanaceae) can be found throughout the world, and there are 14 species, among these, ten species are available in India. Compared with other species, *D. metel*, *D. innoxia*, and *D. stramonium* are the most significant medicinal plants (Schultes and Hofmann, 1979).

D. metel is a perennial plant. The structure of leaves is simple: alternate, dark green, shallowly lobed, and broadly ovate. The appearance of flowers is large, single, and trumpet- growth, extending

branches, and an herbaceous plant of about 1.5 m height. The structure of leaves is simple: alternate, dark green, shallowly lobed, and broadly ovate. The appearance of flowers is large, single, and trumpet-shaped, with a pleasant fragrance and various colors; for instance, in some species, they are white to yellow, and in others, they are light to dark purple. The fruit is egg-shaped, capsulated with short spines, and its diameter is 5 cm. However, it is fond of the worm regions of the world (Drake et al., 1996), as taxonomy shows in Fig. 1.

Under the Solanaceae family, there is evidence of 9–14 species of the *Datura* genus, but not every species has significant distinction to identify it, so only 9 species are widely acceptable; those are *D. ceratocaula* Ortega, *D. discolor* Bernh, *D. ferox* L., *D. innoxia* Mil, *D. kymatocarpa* Barclay, *D. leichhardtii* Benth, *D. metel* L., *D. quercifolia* Kunth, *D. stramonium* L., and *D. wrightii* Regel. *D. metel* is available in Asian and African region (Preissel and Preissel, 2002). Different parts of *D. metel* are shown in Fig. 2.

* Corresponding authors.

E-mail addresses: hdmcoutinho@gmail.com (H.D.M. Coutinho), dmt.islam@bsmrstu.edu.bd (M.T. Islam).

<https://doi.org/10.1016/j.crttox.2023.100106>

Received 4 January 2023; Received in revised form 30 March 2023; Accepted 8 May 2023

Available online 13 May 2023

2666-027X/© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

D. metel contains a lot of significant phytochemicals, including alkaloids, flavonoids, tannins, phenols, cardiac glycosides, amino acids, and carbohydrates (Al-Snafi, 2017b). Various parts of the herb are being utilized as hallucinogenic, narcotic, anti-tussive, antispasmodic, bronchodilator, and anti-asthmatic and are used in epilepsy, diarrhea, hysteria, skin diseases, rheumatic pains, painful menstruation, hemorrhoids, wounds, burns, and skin ulcers. *D. metel* is considered an astringent, bitter, germicide, acrid, anti-phlogistic, anodyne, narcotic, antiseptic, and sedative medicinal plant in Ayurveda (the Ayurvedic Pharmacopoeia of India). Haploid embryos were also developed by Anther culture to produce a haploid *D. metel* plant (Wijesekara and Iqbal, 2021). This review summarizes the phytochemical reports and biological activities of *D. metel* on the basis of database reports.

Results and discussion

Database reports

The latest (till March 20, 2023) searching was performed in the databases: Google Scholar, PubMed, MedLine, Science Direct, and Scopus, with the keywords "Datura metel" and/or paired with "ethno-botanical use", "phytochemicals", and "pharmacological activities". There were no language restrictions. By searching, we have yielded 823 references. From there, 69.99% of the articles were removed for being duplicated throughout the databases; 8.75% were removed for being ineligible by automation tools; 121 articles, or 14.70%, were screened, where 28 articles were excluded for insufficient information; then 76 articles were assessed for eligibility, where 12 articles were excluded for three reasons; and finally, this study included 66 new articles for the review. The PRISMA analysis for data inclusion is shown in Fig. 3.

Traditional uses

Information regarding the traditional uses of natural products and their derivatives has played pivotal roles in the discovery and development of therapeutic agents since ancient times (Pirintsos et al., 2022). Traditionally, *D. metel* is used to treat epilepsy along with skin diseases, hysteria, fever with catarrh, insanity, heart diseases, and diarrhea. For

pain relief, the leaves are used. For treating asthma, this plant is used in China and Vietnam. *D. metel* plant extract is also used as general anesthesia and in the treatment of chronic bronchitis (Ko and Ko, 1999; Kam and Liew, 2002). *D. metel* is also used traditionally in India as a cure for madness or insanity, cerebral complaints and catarrhal infections, elephantiasis, ear discharge, skin diseases, mad dog bites, and indigestion (Tripathi et al., 1996). The flower of *D. metel* is called biamantuoluo in Chinese traditional medicine, where it is mentioned as useful for skin inflammation and psoriasis (Wang et al., 2008). Seeds of *D. metel* are used as an ingredient in tea in Brazil, which provides a sedative effect, and dried flowers of the plant are used for smoking (Monira and Munan, 2012). Young Nigerians also consider *D. metel* as a hallucinogenic agent (Ishola et al., 2021).

In Bangladesh, different parts of the *D. metel* plant are utilized in the treatment of scabies, eczema, and allergies (Ishola et al., 2021), abnormal breathing, pupil problems, pain and swelling, breast pain, asthma, convulsions, and rheumatism (Rahmatullah et al., 2010).

Phytochemical profile

Plant-derived constituents are one of the potential sources of therapeutic bioactives, thus modern medicines. Many important therapeutic medicinal strategies, for example, the basis of therapy of Traditional Chinese medicine (TCM) is composed of natural products and their derivatives, in which phytochemicals are the major candidates (Xiang et al., 2019). Medicinal plants and their isolated compounds are used widely to treat many maladies. Approximately 35% of modern medicine originates from natural sources (Calixto, 2019). *D. metel* has an important place in the traditional systems of medicine as a narcotic, anodyne, and antispasmodic drug similar to Belladonna and Stramonium. Withanolides, tropane alkaloids, and steroidal lactones have been reported from *D. metel*. The plant contains β -sitosterol (Han et al., 2015). *D. metel* also contains many important alkaloids, including litortine, hyoscyamine, acetoxytropine, hyoscyne, fastusine, valtropine, and fastusinine, along with different withanolides, tropine (trigloyl esters), and pseudo-tropine. These are presented in Table 1. However, glycosidase inhibitory activity has been found in the nortropane alkaloids, calystegines, isolated from various species of *D. metel* (Ghani,




Fig. 1. Taxonomy of *Datura metel*.

2003). The root of *D. metel* is rich in atropine, while the aerial parts are rich in scopolamine (Afsharypuor et al., 1995).

Five metelosides A–E and four known compounds were isolated from an acidic methanol whole plant extract of *D. metel*. Cytotoxic effects have been shown for 4, 5, 6, and 2 compounds on MCF-7, SK-Mel-2, and HepG2 cells. On the other hand, three compounds (3, 4, and 7) also moderately showed anti-inflammatory action by suppressing nitric oxide (NO) generation in BV cells, which was stimulated by lipopolysaccharide (LPS)-stimulated (Mai et al., 2017a). Seven glycosides (datunglycosides 1–7) have been isolated from *D. metel* leaves (Tan et al., 2021), as shown in Fig. 4.

Three new compounds (sesquiterpenoid and aliphatic glycoside) have been isolated from *D. metel* roots 70% ethanolic extract, along with 36 known compounds; among these compounds, authors have found that 5 show strong anti-inflammatory activity (Liu et al., 2021). A new compound, phenolic glycoside (methyl 3,4-dihydroxyphenylacetate-4-O-[2-O- β -D-apisoyl-6-O-(2-hydroxybenzoyl)]- β -D-glucopyranoside) is isolated from roots (Qin et al., 2021). Major bioactive compounds are presented in Table 2.

Bioactivities of *Datura metel*

Table 2 reports the diverse biological activities of *D. metel*. Each biological effect has been described in the table.

Antioxidant activity

Oxidative stress leading to inflammatory responses is one of the major defense networks in mammals. However, chronic oxidative stress or inflammation may initiate and progress a variety of diseases (Dandekar et al., 2015). *D. metel* and its derived compounds have been evidently exerting antioxidant (Akharaiyi, 2011; Bhardwaj et al., 2016) and anti-inflammatory effects (Yang et al., 2017) in some studies. A substance having antioxidant and/or anti-inflammatory activity may protect organs, including the nervous system (Islam et al., 2016). The ethanolic and aqueous extractions of the plant showed a significant DPPH radical scavenging capacity between 25.51 and 3.41% and 49.30 and 23.82%, respectively (Akharaiyi, 2011). In this study, a phytochemical report suggests that crude ethanol and water extracts of the different parts of the herb contain alkaloids, saponins, glycosides,

phenols, and flavonoids. The antioxidant capacity of *D. metel* has also been seen by Bhardwaj et al. (2016).

However, in the DPPH model, hydroalcoholic extract of seed showed slightly higher antioxidant action than the methanolic extract, that is hydroalcoholic IC₅₀ value was 25.78 μ g/mL and methanolic extract of seed's IC₅₀ value was 28.34 μ g/mL (Alam et al., 2020; Al-Snafi, 2017a). Importantly, compared with other species like *D. stramonium*, *D. metel*'s antioxidant property is quite high (Iqbal et al., 2017). Antioxidant action is estimated by the aqueous extract of seed and leaf of *D. metel* at 2.5 and 1.5 mg/mL, respectively; they also found that it prolongs the cardiac arrest to 35 to 37 min, when the heart's normal survival time is 14 min (Mbida et al., 2022). Four extract concentrations of four different solvents of *D. metel* extract 25–100 mg/mL were investigated in DPPH scavenging activity, total phenolic content, the reducing power assay, hydroxyl-radical scavenging, and β -carotene bleaching test. The results showed concentration-dependent antioxidant activity; the higher the concentration, the greater the effect (Sangeetha et al., 2014). In another study, the n-hexane extract contained tocopherols, of which 80% were gamma-tocopherol, and this n-hexane extract had the ability to extinguish 40% of the DPPH radical (Ramadan et al., 2007). CeO₂ nanoparticles prepared with *D. metel* ethanolic extract CcO₂ NPs have found DPPH radical scavenging of 16.61% (Yulizar et al., 2020).

Anti-inflammatory and immunomodulatory effects

A unani drug for asthma named Habb-e-Zeequn Nafas (HZN), where *D. metel* is used as the active ingredient, was found to be safer in pre-clinical tests (Firdaus et al., 2022). A newly isolated glycoside from *D. metel* roots (methyl 3,4-dihydroxyphenylacetate-4-O-[2-O- β -D-apisoyl-6-O-(2-hydroxybenzoyl)]- β -D-glucopyranoside) has anti-inflammatory activity (Qin et al., 2021). *D. metel* leaf, seed, and fruit extracts were tested on male albino rats and found to have a nephroprotective effect on kidney function (Imo et al., 2019). Withanolides were found effective against psoriasis, and the mechanism was also established (Cheng et al., 2020). A new drug named 'D. metel L. capsule,' which was approved for treatment and showed more than 90% efficacy and 65% cure rates (Su et al., 2022). Successful Cu₂O nanoparticles were prepared by using *D. metel* extract, but their effect was not determined (Chinnaiah et al.,




Fig. 2. Different parts of *Datura metel* [(a) plant, (b) flower, (c) unripe green fruits, (d) ripe canary fruits, (e) mature black-seeds, (f) roots of *D. metel*].




Fig. 3. PRISMA analysis of database reports for *Datura metel*.

Table 1
Constituents of alkaloids of *Datura metel*.

<i>D. metel</i> parts	Alkaloidal presence (%)	Major constituents	Sources
Leaves	0.426%	Hyoscyamine, scopolamine and atropine, withanolides	Siddiqui et al., 1987; Dabur et al., 2005
Roots	0.35%	3 α -tigloyloxytropine, hyoscyne, hyoscyamine, norhyoscyne, tropine, apohyoscyne, tigloidine and cuscohygrine	Ghani, 2003
Seeds	0.426%	Daturanalone, hyoscyamine and fastusic acid and some tropane alkaloids	
Fruits (Pericarp)	Not quantified	Daturanalone, β -sitosterol, daturadiol, and tri-terpene	Liu et al., 2020
Seeds	Not quantified	4 new indole alkaloids daturametelindoles A-D (1–4)	De, 2003
Propagated shoots	Not quantified	withanolide-12-deoxywithastramonolide and C28 sterol 3 β ,24 ξ -dihydroxy-ergosta-5, 25-dienolide compounds	
Cultured callus	Not quantified	5 α -pregnane3 β ,20 β -diol and Cholesterol compound	
Flowers	Not quantified	5 compounds of withanolide (baimantuoluoline A-C and withametelin C and withafastuosin E), withametelins 10 compounds I-P, and 12 β -hydroxy-1,10-secowithametelin B and 1,10secowithametelin B compounds.	Agharkar, 1991; Manickam et al., 1993; Yang et al., 2010a

2022).

Anti-microbial activity

Anti-bacterial effect

Anti-bacterial activity is a very intricate process that involves microscopic creatures whose ability to reproduce and function at every stage of life, including nourishment, metabolism, and respiration, can be impacted by the presence of potentially harmful elements (Patachia and Croitoru, 2016). A zone of inhibition is a region of media in which bacteria are unable to grow due to the presence of a medication that inhibits their growth. The lowest inhibitory concentration of an anti-microbial medication that precludes observable growth of a bacterium after overnight incubation with medium (Barnard, 2019).

The ethanolic and aqueous extracts (20 mg/mL) of leaf, stem bark and roots of *Datura* acted against, *Streptococcus dysenteriae* (gram-negative), *Pseudomonas aeruginosa* (gram-negative), *Bacillus cereus* (gram-positive), *Escherichia coli* (gram-negative), *Klebsiella pneumoniae* (gram-negative), *Staphylococcus aureus* (gram-positive), and *Streptococcus β hemolytic* (gram-positive) (Akharaiyi, 2011). In this study, the leaves and stem bark extracts produced zones of inhibition between 12 and 35 mm, while the aqueous extract by 10 to 22 mm, and by ethanol extract 12 to 32 mm. The MBC was 10 to 20 mg/mL. So, *D. metel* has inhibitory action against both of the gram-positive and gram-negative strain and *S. aureus* was the mostly suppressed bacteria by the ethanolic extract.

According to Bachheti et al. (2018), at least 7 bacterial strains showed the highest inhibition zone and lowest MIC when treated with the seed oil, including *Pseudomonas aeruginosa* (18 mm) and *Lactobacillus delbrueckii lactis* (19 mm). Studies have shown that the higher the oil concentration, the greater the antibacterial response. Methanolic extract depicted ($p < 0.05$) antibacterial effect and biofilm inhibition against *Bacillus subtilis*, MRSA, and *E. coli* at 94, 88, and 92%, respectively (Prasathkumar et al., 2022). Silver nanoparticles (25–70 nm) were




Fig. 4. Newly isolated seven daturglycosides (1–7).

prepared in *D. metel* extract with ultrasound and found to have a higher zone of inhibition on tested bacteria (Nethradevi et al., 2012; Fatimah et al., 2021).

Anti-fungal effect

In a study, the chloroform, hexane, methanolic, and acetone fractions of *D. metel* were found to act against *Aspergillus fumigatus*, *A. niger*, and *A. flavus* (Rajesh and Sharma, 2002). The minimum inhibitory concentration found for the chloroform solvent was 625.0 µg/mL. A derivative of pyrrole, named- 2β-(3,4-dimethyl-2,5-dihydro-1H-pyrrol-2-yl)-10-methylethyl pentanoate, separated from the leaf of the plant, was found to act against *A. niger*, *Candida albicans*, *A. flavus*, *C. tropicalis*, and *Aspergillus fumigatus* (MIC value: 87.5 mg/mL) (Dabur et al., 2004). In a disc diffusion assay, the MIC was observed at 12.5 mg disc, and the methanol extract of *D. metel* was investigated against the *Aspergilli*, and the MIC was observed 1.25–2.50 mg/ml (Dabur et al., 2004; Monira and Munan, 2012). *D. metel* extract of leaves and stem showed antifungal properties; these authors used 6 methanol concentrations (1, 1.5, 2, 2.5, 3, and 3.5%), and found leaf 3.5% conc. has ability to inhibit *Rizoctonia solani* Kuhn at 75%, which is greater than stem at same concentration (Hanif et al., 2022).

Anti-viral effect

An atropine isolated from the *Datura* inhibited the growth of enveloped viruses (e.g., HBV, HCV, HIV, and influenza viruses) and also inhibited the glycosylation of viral proteins of the Herpes virus, resulting in no formation of virions (Yamazaki and Tagaya, 1980). Effective against tomato mosaic virus (TMV), activated oxygen species play the defensive role against the virus and program cell death to save the plant from this virus (Al-Huqail and Aref, 2017).

Wound healing activity

The process of returning damaged tissue's tissue structure to its pre-injury state as closely as feasible is known as wound healing. Its characteristics include coagulation factors, inflammatory processes, re-epithelialization, granulation, modification of the extracellular matrix, and scarring (Boakye et al., 2018). It is reported that the seed of *D. metel* is externally used for piles (Yusuf et al., 1994; Yusuf et al., 2009). *Datura* ointment at 10% w/w, in an excision wound model, exhibited significant wound healing capacity in Wistar albino rats. The result was comparable to the standard, nitrofurazone 0.2% w/w, which was used in this study (Vimal and Suseela, 2010).

Studies have found that pre-ingestion of seed extract of *Datura* increased the viable rate of the rat after toxication with dichlorvos; the possible reason could be the availability of a significant percent of atropine in seeds and other anticholinergic compounds in it (Das et al., 2012; Manpreet et al., 2017). Providing *D. metel* seed extract 7.5 mg/kg intraperitoneal injection to male rats before giving 25 mg/kg dichlorvos inj. showed longer times of survival, even 24 h more survival in the case of *Datura*-treated animals (Banja et al., 2004; Al-Snafi, 2017a). The wound-healing capacity of *D. metel* and its isolated compounds confirm the medicinal value of this plant (Vimal and Suseela, 2010).

Herbicidal activity

A weed-killer or herbicidal compound is a substance used to kill plants, particularly weeds or other undesirable species (Almarie, 2022). The n-hexane, methanol, and water extracts of root and shoot at 5, 10, and 15% (w/v) of the plant exerted noticeable herbicidal action on *Phalaris minor* Retz. Moreover, the methanol (5–15%) and n-hexane (15%) root extracts of this plant also considerably lowered the germination, shoot, and root lengths of the sample (Javaid et al., 2008).

Insecticidal activity

The methanolic extract of *Datura* seeds at 2.5, 5.0, 7.5, and 10.0% were found to act against *Helicoverpa armigera* (Hubner) (Singh and Singh, 2008). This study was also supported by the authors Monira and Munan (2012) and Al-Snafi (2017a), who said methanolic extract of *D. metel* seeds significantly blocked different developmental stages such as larval, pupal, pupation quantity, and adult emergence.

Anti-cancer activity

Different glycosidases are effectively inhibited by nitrogen-rich polyhydroxylated compounds. They can be utilized for treating many diseases, including cancer, diabetes, immunological diseases, and virus-causing infections (Wang et al., 2019). Including *Datura*, other 14 Solanaceae family plants contain a significant number of withanolides, and these compounds have a notable anti-cancer effect by creating oxidative stress and downregulating NF-κB, transcription factor STAT3, and Hsp90. The inhibition of PI3K/Akt and MAPK pathways results in cytoskeletal and structural protein dysregulation. Cancer cell angiogenesis is inhibited via HIF-1 downregulation (Akhtar et al., 2020; Nkwe et al., 2021), a possible anticancer pathway depicted in Fig. 5. Withanolides have lots of significant biological activities, for example, anti-tumor activity (Bellila et al., 2011), and importantly, anti-

Table 2
Bioactivities of phytoconstituents of *Datura metel*.

Compounds name	Parts/derivatives	Extraction process	Result/activity	References
Withafastuosin	Flower	70% Ethanol used	Depicted activity for psoriasis	Yang et al., 2007
Two new lignin compounds, Compound 1 (C ₂₁ H ₂₂ O ₇), compound 2 (C ₂₇ H ₃₂ O ₁₂) and known 18 lignin compounds	Roots	70% Ethanol	Cytotoxicity test against Hela cell line IC ₅₀ for compound 1 (48.20 μM) and for compound 2 (58.86 μM)	Yang et al., 2018
Withametelins	Flower	Methanolic extraction	Cytotoxicity showed	Pan et al., 2007
Sesquiterpenoids, nine new (1–9) and three known (10–12)	Leaves	Ethanol extract ethanol–water (7:3, v/v)	Anti-inflammatory activity against RAW264.7 and IC ₅₀ value was 9.33–11.67 μM, control group was greater thana this.	Tan et al., 2020a
Withanolides	Flower	Used 50% Ethanol	Presented activity for psoriasis	Yang et al., 2007
Ophiobolin A	Produced by its endophytic fungus (<i>Bipolaris</i> sp.)	–	Anti-cancer activity	Maehara et al., 2020
Baimantuoluoline A	Flower	50% Ethanol	Showed the activity for psoriasis	Yang et al., 2007
(5α,6β,7α,22R)-5,6,7,27-tetrahydroxy-1-oxowitha-2,24-dien-27-O-β-D-glucopyranoside	Flower	Extracted with 70% Ethanol	For treating of psoriasis	Yang et al. (2010b)
6,7-dimethyl-1-D-ribityl-quinoxaline-2,3(1H,4H)-dione-5'-O-β-Dglucopyranoside	Flower	Used 70% Ethanol	Psoriasis treatment	Yang et al. (2010b)
Sesquiterpenoid and aliphatic glycoside (E)-methyl 4-(3-(4hydroxyphenyl)-N-methylacrylamido)butanoate	Roots Flower	70% Ethanol 70% Ethanol used for extraction	Anti-inflammatory activity by 5 compounds Treatment of psoriasis	Liu et al., 2022 Yang et al., 2010a
Ergostane-type C28 sterols, daturmetesides A-E (1–5)	Leaves	70% Ethanol	Anti-inflammatory activity, IC ₅₀ values ranging from 17.05 ± 0.35 μM to 24.88 ± 0.93.	Tan et al., 2020b
1, 10-seco-withametelin B	Flower	Methanolic extraction	Cytotoxicity is observed	Pan et al., 2007
12 β -hydroxy-1,10-seco-withametelin B	Flower	Methanolic extraction	Cytotoxicity is observed	Pan et al., 2007
Alkaloid datumetine	Leaves	Para-methoxy benzoic acid	Anti-spasmodic action	Siddiqui et al., 1987
Lignin-amides from <i>Datura metel</i> seeds (LDS)	Seeds	95% Ethanol	Anti-neuroinflammatory, Alzheimer disorder	Wang et al., 2021
2-(3,4-dimethyl-2,5-dihydro-1H-pyrrol-2-yl)-1-methylethyl pentanoate	Leaves	Chloroform	Antifungal activity, MIC 87.5 μg/ml	Dabur et al., 2005
Melatonin	Flower	Methanolic extract	Cold stress controlling	Murch et al., 2009
Serotonin	Flower	Methanolic extract	Induced during stress-condition	Murch et al., 2009
2 new compounds- meteloside F(1) and meteloside G(2) and 6 known steroids	Seeds	95% Ethanol	Inhibiting nitric oxide production from RAW 264.7 cells, IC ₅₀ value 30.2–44.8 μM	Yang et al., 2020
Seven new glycoside and seventeen known	Leaves	70% Aqueous ethanol	Antiproliferative activity to MGC-803, Hela, and Ishikawa cell-lines	Tan et al., 2021
A phenolic glycoside name- methyl 3,4-dihydroxyphenylacetate-4-O-[2-O-β-D-apisoyl-6-O-(2-hydroxybenzoyl)]-β-D-glucopyranoside	Roots	Ethanol	Anti-inflammatory activity	Qin et al., 2021
Found Alkaloid (88.77 ± 1.01 mg AE/g), Phenolic (124.61 ± 0.68 mg GAE/g), tannins (38.72 ± 0.51 mg GAE/g), flavonoids (42.24 ± 0.18 mg QE/g)	Leaves	Methanol	Antibacterial, anti-diabetic, antioxidant, anti-inflammatory, cytotoxic effects etc.	Prasathkumar et al., 2022
13 new withanolide aglycones, baimantuoluolines L-X, with baimantuoluoside J (a new withanolide glycoside)	Flowers	Ethanol	Immunosuppressive by 1–14 compounds and anti-proliferative action against SGC-7901, HepG2, MCF-7 were tested.	Liu et al. (2020a)
12 undescribed withanolides	Leaves	Ethanol	Inhibits nitric oxide production, IC ₅₀ value for daturafoliside L, T, M ranging from 9.37 to 18.64 μM and for K, R, A range 22.84–33.36 μM	Guo et al., 2018
2 guanine sesquiterpenes, (1)1β,5α,7β-guaiene-4β,10α,11-triol and (2)1α,5α,7α-11-guaiene-2α,3β,4α,10α,13-pentaol, along with 8 described compounds	Whole plant	Methanol	Anti-inflammatory effect was ensured by NO inhibition.	Mai et al., 2017
Daturaolone	Fruits	Chloroform	Potent anti-fungal and antibacterial action. Effective against <i>B. subtilis</i> , <i>Klebsiella pneumonia</i> , <i>S. epidemidis</i> , <i>S. aureus</i> and zone of inhibition found 12–30 mm. Moreover, fruit extract provide modest sensitivity	Bawazeer, and Rauf, 2021

(continued on next page)

Table 2 (continued)

Compounds name	Parts/derivatives	Extraction process	Result/activity	References
Daturaturin A (DTA)	Dried Flowers of <i>D. metel</i>	Ethanol	against <i>C. glabrata</i> , <i>T. longifusus</i> , <i>F. solani</i> , <i>C. albicans</i> , <i>M. canis</i> , <i>A. flavus</i> Anti-inflammatory and anti-proliferative effect and anti-psoriasis by activation of autophagy.	Wei et al., 2021
6 known terpenoids (2–6) and a new-diterpenoid called-kaurane daturaside A(1)	Dried pericarps	Ethanol	Compounds 2 and 5 showed anti-inflammatory action against LPS induced RAW264.7 cells (IC ₅₀ value is lower than 11 μM).	Liu et al., 2021a
12 phyto constituents including (tetradecanoic acids, eugenol, dodecanoic, 2-pentadecanone 6,10,14 trimethyl, 9, 12 octadecanoic acid, pentadecanoic acid, pentadecanoic acid, 6-octadecanoic acid, 1 4-methyl- methyl ester, n-hexadecanoic, 9,12,15-octadecatrienoic acid, phytol and heptacosane)	Leaves and stem	Methanol	Antifungal effects	Hanif et al., 2022
A total of ten substances, including scoptone, isofraxidin, daturadiol, arenarine D, 1–4-benzenediol, vanillin, G-sitosterol, N-trans-feruloyl-tyramine, scopoletin, and hyoscyamilactol	Seed extract	–	–	Han et al., 2015




Fig. 5. Possible anticancer mechanism of withanolides (Here withanolides increase oxidative stress and downregulating nuclear factor kappa light chain B (NF-kB), transcription factor signal transducer and activator of transcription 3 (STAT3), heat-shock protein 90 (Hsp90) via phosphoinositide 3-kinase (PI3K)/protein kinase B (Akt) and mitogen-activated protein kinase (MAPK) pathways which results cytoskeletal and structural protein dysregulation; thus, decrease angiogenesis and increase cytotoxic activity against cancer cell).

inflammatory and immunomodulatory effects (Yu et al., 2017). The withanolides have inhibitory effects on tumor cell proliferation, and they can act as angiostatic agents. Moreover, they may induce the phase-II enzyme quinone reductase (Pan et al., 2007).

Datura is evidently a source of calystegines. Some withanolide glycosides (aturametelins), daturaturin, 7,27-dihydroxy-1-oxo and 2,5,24-trienolide are also separated from the methanol extraction process of *D. metel* aerial parts. The non-glycosidic compound of the plant exhibited a significant antiproliferative effect on the HCT-116 cell line (IC₅₀: 3.2 ± 0.2 μM) (Monira and Munan, 2012). In another study, daturanolides A–C, along with 6 previously described withanolides (4–9), were found in the flowers of *D. metel* (Wang et al., 2019), as shown in Fig. 6. In this study, compound 6 exhibited marked cytotoxic effects on five human cancer cell lines, such as HepG2, NCI-H460, HCT116, BGC823, and U87-MG. When NF-kB is inhibited, the genes are downregulated, which prevents cancer cells from proliferating and forming new blood vessels. Apoptosis induction in cancer cells may be aided by blocking the NF-kB signaling pathway as well as TNF-related apoptosis-inducing ligands (Varma et al., 2011).

The sesquiterpenoid glycosides, citroside A, dmetelisproside A, and staphylionoside D, separated from the *D. metel* leaves (Fig. 6), were evaluated for cytotoxic effects towards MDAMB-231, SGC-7901, HeLa, Hepg2, and MCF-7 cancer cell lines, and anti-inflammatory activity was observed on the RAW 264.7 cells. Dmetelisproside A and citroside A showed strong cytotoxicity against HeLa and SGC-7901 cells (IC₅₀: 21.43–29.51 μM). Staphylionoside D, citroside A, dmetelisproside A, and citroside A also inhibited nitric oxide production, with IC₅₀ values of 44.31, 34.25, and 31.10 μM, respectively (Guo et al., 2019). Another study have found 3 newly glycosides named- daturametelins H to J they have tested with other two established compound daturaturin-A and 7, 27-dihydroxy-1-oxowitha-2,5,24-trienolide against HCT-116 (human colorectal carcinoma) and have found the IC₅₀ values 3.2 ± 0.2 μM.

According to the studies of Al-Snaifi (2017b) and Ibrahim et al. (2018), the stem and root extracts of *D. metel* have demonstrated cytotoxic actions against HepG-2 (human liver cancer cells), where IC₅₀ values were 341.12 and 613.88 μg/mL, respectively. On the other hand, the root and leaf extracts were effective against HeLa (cervical carcinoma cells) with IC₅₀ values of 348.35 and 267.76 μg/mL, respectively.




Fig. 6. Chemical structure of some potential anticancer compounds of *Datura metel*.

The IC_{50} of methanolic extraction of *D. metel* fruits towards the Vero-cell line was found to be 3 mg/mL; in this study, the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) method was utilized in an in vitro cytotoxicity assay in the Vero-cell line (Roy et al., 2016). Tan et al. (2020a) have evaluated 24 compounds of *D. metel* against Hela, Ishikawa, and MGC-803 cell lines and found seven compounds provided antiproliferative action.

Hypoglycemic effect

Hypoglycemia is the state of having blood sugar (glucose) levels that are below normal. The body uses glucose as its main source of energy. Diabetes treatment frequently involves managing hypoglycemia (Wild et al., 2007). *D. metel* seed extract showed hypoglycemic action in alloxan-stimulated diabetic rats (Krishna Murthy et al., 2004). The authors demonstrated that a rapid normalization of blood glucose levels was also seen, possibly due to the increase in protection and secretion of insulin's capabilities by the β -cells.

Neuroprotective effects

Neuroprotection means the processes and tactics used to guard the central nervous system (CNS) against injury caused by both acute (e.g., trauma or stroke) and chronic (e.g., dementia, Parkinson's, Alzheimer's, seizures, etc.) neurodegenerative illnesses (Rehman et al., 2019). In a

study, a methanolic crude extract of *D. metel* at 0.6, 1.2, 1.5, 2, and 2.4 g/kg demonstrated a dose-dependent oral anesthesia in dogs (Babalola et al., 2013). The aqueous extracts of seed and leaf of *D. metel* are also evident to show neuropsychopharmacological effects in rat and mouse models (Abena and Miguel, 2004). In a study, the aqua extract of *D. metel* seed at 400 and 800 mg/kg was found to increase neuron-motor activity while lowering the duration of barbituric acid-induced sleeping time, antagonizing catalepsy and inducing ptosis by haloperidol, and inducing immobility in the test animals (Al-Snafi, 2017a). Importantly, NMDAR is a glutamate receptor and ion channel that can be noticed in neurons and is affected by datumetine (the active compound of *D. metel*) and results in intoxication and memory deficit (Ishola et al., 2021).

Anti-nociceptive effect

The drugs known as analgesics are used to treat pain. Analgesics don't cut off nerves, impair your ability to detect your surroundings, or affect consciousness, unlike the drugs used as anesthetics during surgery. They are referred to as painkillers or pain relievers occasionally (Price, 1999). The aqueous seed extraction of *Datura metel* did not exhibit analgesic effects on experimental animals (Wannang et al., 2009). However, the alcoholic seed extract exerted a dose-dependent analgesic effect; the ED_{50} was 25 for aqueous seed extraction and 50

mg/kg for the alcoholic extract in the hot plate and formalin tests applied on rats (Khalili and Atyabi, 2004). In a study indomethacin was used as standard drug and compared the result of ethanolic root extract of *D. metel* in rat and result showed anti-inflammatory activity at 200 mg/kg, comparing Indomethacin (10 mg/kg). On the other hand, an aquas extract of *D. metel* seeds and leaves was found effective at 800 and 400 mg/kg dosage in mice, and inflammation was induced by acetic acid (Al-Snafi, 2017).

Daturaolone is a potential compound found in *D. metel*; in a study, it was found that this compound can inhibit cyclooxygenase and lipoxygenase to provide anti-inflammatory and analgesic activity (Rauf et al., 2016); the basic mechanism is to inhibit cyclooxygenase COX-1 and COX-2 and lipoxygenase; this inhibition causes the suppression of prostaglandin synthesis or might be the narcotic effects of *Datura* species (Chandan et al., 2021; Jaafar et al., 2018). In the process of inflammation, arachidonic acid is crucial because it serves as a building block for the creation of two different types of vital compounds, prostaglandins and leukotrienes, in the body, and daturaolone inhibits the production of prostaglandins and leukotrienes by cyclooxygenase 1 and 2 and lipoxygenase inhibition (Samuelsson, 1987; Baig et al., 2021), as shown in Fig. 7.

Contraceptive effects

Antifertility medications are chemical compounds that block the activity of hormones that encourage pregnancy. These medicines serve as a form of protection and actually lower the likelihood of pregnancy (Rivera et al., 1999). Usually, antifertility medications are derived from synthetic progesterone derivatives or a mix of progesterone and estrogen derivatives (Druckmann, 2009). Plant sources can act as contraceptives or antifertility agents (Brondegaard, 1973) Contraceptive action of *D. metel* is considerably observed in albino mouse trial by the acetone extracts of seeds, where three concentrations were used (0.5%, 1% and 2%) orally given to female albino. These mice were given the opportunity to mate after 15 days of treatment with normal male albinos. Dissection was performed after ten days and it was found that 2% seed extract showed full (100%) anti-fertilization, whereas 0.5% and 1% seed extracts presented 80% and 40% anti-pregnant action, respectively. This author has commented that *D. metel* is a good source of contraceptive compounds with inconsiderable side effects (Pandiarajan et al., 2012). Not only this, there was another study on male rats where *D. fastuosa*'s (*metel*) ethanolic extract was given orally for 7 weeks at 2, 4, and 6 mg/kg concentration, and the results were surprising as it decreased sperm content and normal sperm compared with the control group. Moreover, hormonal levels of testosterone, luteinizing hormone (LH), follicle-stimulating hormone (FSH), and testicle weight also decreased; as a result, pregnancy occurrence also decreased considerably (Al-Mailay, 2008). The antifertility activity of *D. metel* is shown in Fig. 7. However, in these studies, there was no specific identification of a compound that could be active as a contraceptive.

Metabolism of *Datura metel* seed compounds

We have found a study where authors used 0.15 mg/kg oral *D. metel* seeds aquas extract in rats to find out the metabolism of *D. metel* compounds. They have characterized 42 compounds from seed extract and studied thirteen bioactive compounds metabolic pathways. They found 61 compounds in blood plasma, 81 in urinary excretion, and 76 in stools. Though this might be the first study of full metabolic profiling of *D. metel* seeds, proper quantification is not mentioned (Xia et al., 2019).

Toxicological profile of *Datura metel*

More research on poisonous plants is crucial to avoid livestock losses (Panter et al., 2013). It is due to the fact that each plant contains thousands of compounds, and among them all, some may not be poisonous to us (Dutkiewicz et al., 2016). In fact, the toxic substances can be used for many purposes, such as herbicidal (Javaid et al., 2010), insecticidal, pesticidal, to kill cancer cells, and so on (Tariq et al., 2015; Shuping and Eloff, 2017). *Datura* is a toxic medicinal plant. All the parts of this herb are poisonous. It is because of the presence of poisonous tropane alkaloids and anticholinergic agents (e.g., atropine, hyoscyamine, and scopolamine). Convulsions, acute confusion, dry mouth, fever, tachycardia, urine retention, hot-flushed dry skin, hallucinations, headache, dilated pupils, delirium, a weak and rapid pulse, coma, and even death are common signs and symptoms of *Datura* poisoning (Kam and Liew, 2002). The withametelins I, K, L, and N showed cytotoxicity activities against K562 (leukemia), BGC-823 (gastric), and A549 (lung) cancer cell lines, with IC₅₀ values of 0.05 to 3.5 μM. Withamelin J depicted moderate cytotoxic activity against K562 and BGC-823 but less cytotoxicity against the A549 cell line (Pan et al., 2007). Other cytotoxic withametelins are J, M, O, P, 12-β-hydroxy-1,10-seco-withametelin B, and 1,10-seco-withametelin B.

Importantly, *D. metel* exhibited safety up to a 2000 mg/kg body weight dosage since it creates no observable sign of toxicity or death. Histological studies showed the minimized weight of the organs and necrotic modifications in the liver, followed by the increasing of ALP (serum alkaline phosphatase), SGOT (serum glutamic-oxaloacetic transaminase), and GPT (glutamyl pyruvic transaminase) (Bouzidi et al., 2011, Ogunmoyole et al., 2019).

Tropane alkaloids can be found throughout the plant, particularly in the leaf and seeds, and are poisonous. It includes atropine, hyoscyamine, and scopolamine. The central and peripheral muscarinic neurotransmission is inhibited by the tropane alkaloids, resulting in the anticholinergic syndrome (Schultes and Hofmann, 1979; Wijesekara and Iqbal, 2021; Kam and Liew, 2002); a possible mechanism is shown in Fig. 8.

Conclusion and future perspectives

D. metel possesses several phytochemicals with a wide range of pharmacological effects. Despite its toxicity, the plant holds significant promise as a potential source of plant-based therapeutics. Future research should focus on exploring the plant's therapeutic potential while ensuring proper safety precautions are taken. Advancements in




Fig. 7. Proposed anti-inflammatory and analgesic effect of daturaolone (Through COX-1 and COX-2 and lipoxygenase inhibition it may inhibits prostaglandins and leukotrienes production thus reduce in inflammation).




Fig. 8. Anti-fertility activity of *Datura metel* (Here, ethanolic extract of *D. metel* showed sperm content lowering activity this might be the result of bioactive compound of this plant decreasing testosterone, luteinizing hormone (LH) and follicle-stimulating hormone (FSH). On the other hand, acetone seed extract on female mice showed 100 percent antifertility action with 2% extract concentration).




Fig. 9. Proposed toxicity mechanism of tropane alkaloids of *Datura metel* (Here, tropane alkaloids blockade the function of nicotinic and muscarinic receptors and neurotransmitter like acetylcholine (ACh) cannot bind with the targeted receptor which results anticholinergic syndrome).

technology and methodologies could lead to the discovery of novel compounds with beneficial biological activities that could help treat various diseases. *D. metel*, like many other medicinal plants, has the potential to contribute to the development of modern medicine and improve the quality of life for millions of people. Therefore, further exploration of the plant's pharmacological properties could lead to the development of new and effective drugs for the treatment of various ailments (Fig. 9).

A number of alkaloids found in the plant, such as atropine, scopolamine, and hyoscyamine, have been demonstrated to have anticholinergic, antispasmodic, and analgesic properties. *D. metel* is a viable option for the treatment of neurological conditions, including Alzheimer's and Parkinson's disease, because recent research has also indicated that it may have neuroprotective and anti-inflammatory qualities. Nonetheless, due to the plant's recognized toxicity and misuse potential, use for medical purposes must be done with caution. Future treatments and therapies will likely be more successful as a result of new insights into the possible advantages and hazards of *D. metel* that are anticipated to be revealed as research into the plant progresses.

Funding

This research received no external funding.

Data availability statement

Not applicable.

CRedit authorship contribution statement

Tawhida Islam: Conceptualization. **Iffat Ara:** Methodology. **Tariqul Islam:** Conceptualization. **Pankaj Kumar Sah:** Methodology. **Ray Silva de Almeida:** Software. **Edinaldo Fagner Ferreira Matias:** Resources. **Cícero Lucas Gomes Ramalho:** Resources. **Henrique Douglas M. Coutinho:** Writing – original draft, Project administration. **Muhammad Torequl Islam:** Conceptualization, Supervision, Project administration.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

References

- Abena, A.A., Miguel, L.M., 2004. Neuropsychopharmacological effects of leaves and seeds extracts of *Datura fastuosa*. *Biotechnology* 3, 109–113.
- Afsharypuor, S., Mostajeran, A., Mokhtary, R., 1995. Variation of scopolamine and atropine in different parts of *Datura metel* during development. *Planta Med.* 61, 383–384.
- Agharkar, S.P., 1991. *Medicinal Plants of Bombay Presidency*. Scientific Publishers, Jodhpur.
- Akharaiyi, F.C., 2011. Antibacterial, Phytochemical and Antioxidant activities of *Datura metel*. *Int. J. Pharm. Tech. Res.* 3, 478–483.
- Akhtar, N., Baig, M.W., Haq, I.U., Rajeev, V., Cutillas, P.R., 2020. Withanolide metabolites inhibit PI3K/AKT and MAPK pro-survival pathways and induce apoptosis in acute myeloid leukemia cells. *Biomedicine* 8 (9), 333.
- Alam, W., Khan, H., Khan, S.A., Nazir, S., Akkol, E.K., 2020. *Datura metel*: a review on chemical constituents, traditional uses and pharmacological activities. *Curr. Pharm. Des.* 27, 2545–2557.
- Al-Huqail, A.A., Aref, N.M.A., 2017. Physiological parameters correlated with Tomato Mosaic Virus inducing defensive response in *Datura metel*. *Saudi J. Biol. Sci.* 24 (4), 864–874.
- Al-Mailay, H.K.A., 2008. The effect *Datura fastuosa* L. alcohol extract on the fertility of white rats males. *J. Al-Qadisiyah Pure Sci.* 13 (3), 1–11.

- Almarie, A.A., 2022. Phytotoxic Activity of Essential Oils. In: *Essential Oils: Applications and Trends in Food Science and Technology*. Springer International Publishing, Cham, pp. 263–279.
- Al-Snafi, A.E., 2017a. A review on *Dodonaea viscosa*: a potential medicinal plant. *IOSR J. Pharm.* 7 (2), 10–21.
- Al-Snafi, A.E., 2017b. Medical importance of *Datura fastuosa* (syn: *Datura metel*) and *Datura stramonium*-a review. *IOSR J. Pharm.* 7, 43–58.
- Babalola, S.A., Sulaiman, M.M., Hassan, A.Z., Adawa, D.A.Y., 2013. Evaluation of the crude methanolic seed extract of *datura metel* L. as a potential oral anaesthetic in dogs. *Veterinary Res.* 6, 115–119.
- Bachheti, R.K., Rai, I., Mishra, V.K., Joshi, A., 2018. Antioxidant and antimicrobial properties of seed oil of *Datura metel*. *J. Environ. Biol.* 39 (2), 182–188.
- Baig, M.W., Fatima, H., Akhtar, N., Hussain, H., Okla, M.K., Al-Hashimi, A., Al-Qahtani, W.H., AbdElgawad, H., Haq, I.U., 2021. Anti-inflammatory potential of *Daturaolone* from *Datura innoxia* Mill.: in silico, in vitro and in vivo studies. *Pharmaceutic* 14 (12), 1248.
- Bania, T.C., Chu, J., Bailes, D., O'Neill, M., 2004. Jimson weed extract as a protective agent in severe organophosphate toxicity. *Acad. Emerg. Med.* 11, 335–338.
- Barnard, R.T., 2019. The zone of inhibition. *Clin. Chem.* 65 (6), 819.
- Bawazeer, S., Rauf, A., 2021. In vitro antibacterial and antifungal potential of amyrin-type triterpenoid isolated from *datura metel* Linnaeus. *Biomed Res. Int.* 2021.
- Bellila, A., Tremblay, C., Pichette, A., Marzouk, B., Mshvildadze, V., Lavoie, S., Legault, J., 2011. Cytotoxic activity of withanolides isolated from Tunisian *Datura metel* L. *Phytochemistry* 72, 2031–2036.
- Bhardwaj, K., Kumar, S., Ojha, S., 2016. Antioxidant activity and ft-ir analysis of *Datura innoxia* and *Datura metel* leaf and seed methanolic extracts. *Afric. J. Tradit. Complement. Altern. Med.* 13, 7–16.
- Boakye, Y.D., Agyare, C., Ayande, G.P., Titiloye, N., Asiamah, E.A., Danquah, K.O., 2018. Assessment of wound-healing properties of medicinal plants: The case of *Phyllanthus muellerianus*. *Front. Pharmacol.* 9, 945.
- Bouzidi, A., Mahdeb, N., Kara, N., 2011. Toxicity studies of alkaloids of seeds of *Datura stramonium* and synthesis alkaloids in male rats. *J. Med. Plants Res.* 5 (15), 3421–3431.
- Brondegaard, V.J., 1973. Contraceptive plant drugs. *Planta Med.* 23 (02), 167–172.
- Calixto, J.B., 2019. The role of natural products in modern drug discovery. *An. Acad. Bras. Cienc.* 91, e20190105.
- Chandan, G., Kumar, C., Chibber, P., Kumar, A., Singh, G., Satti, N.K., Gulilat, H., Saini, A.K., Bishayee, A., Saini, R.V., 2021. Evaluation of analgesic and anti-inflammatory activities and molecular docking analysis of steroidal lactones from *datura stramonium* L. *Phytomedicine* 89, 153621.
- Cheng, Y., Liu, Y., Tan, J., Sun, Y., Guan, W., Jiang, P., Yang, B., Kuang, H., 2020. Integrated serum metabolomics and network pharmacology approach to reveal the potential mechanisms of withanolides from the leaves of *Datura metel* L. on psoriasis. *J. Pharm. Biomed.* 186, 113277.
- Chinnaiah, K., Maik, V., Kannan, K., Potemkin, V., Grishina, M., Gohulkumar, M., Tiwari, R., Gurushankar, K., 2022. Experimental and theoretical studies of green synthesized Cu2O nanoparticles using *Datura Metel* L. *J. Fluoresc.* 32 (2), 559–568.
- Dabur, R., Ali, M., Singh, H., Gupta, J., Sharma, G.L., 2004. A novel antifungal pyrrole derivative from *Datura metel* leaves. *Die Pharmazie –Int. J. Pharm. Sci.* 59, 568–570.
- Dabur, R., Chhillar, A.K., Yadav, V., Kamal, P.K., Gupta, J., Sharma, G.L., 2005. In vitro antifungal activity of 2-(3,4-dimethyl-2,5-dihydro-1H-pyrrol-2-yl)-1-methylethyl pentanoate, a dihydropyrrole derivative. *J. Med. Microbiol.* 54, 549–552.
- Dandekar, A., Mendez, R., Zhang, K., 2015. Cross talk between ER stress, oxidative stress, and inflammation in health and disease. *Meth. Mol. Biol.* 1292, 205–214.
- Das, S., Kumar, P., Basu, S.P., 2012. Phytoconstituents and therapeutic potentials of *datura stramonium* Linn. *J. Drug Deliv. Ther.* 2, 4–7.
- De, B., 2003. Steroidal compounds from in vitro regenerated shoots of *Datura metel*. *Fitoterapia* 74, 14–17.
- Drake, L.R., Lin, S., Rayson, G.D., Jackson, P., 1996. Chemical modification and metal binding studies of *Datura innoxia*. *Environ. Sci. Technol.* 30, 110–114.
- Druckmann, R., 2009. Profile of the progesterone derivative chlormadinone acetate—pharmacodynamic properties and therapeutic applications. *Contraception* 79 (4), 272–281.
- Dutkiewicz, J., Mackiewicz, B., Lemieszek, M.K., Golec, M., Milanowski, J., 2016. *Pantoea agglomerans*: a mysterious bacterium of evil and good. Part IV. Beneficial effects. *Ann. Agric. Environ. Med.* 23, 206–222.
- Fatimah, I., Hidayat, H., Nugroho, B., Husein, S., 2021. Green Synthesis of silver nanoparticles using *datura metel* flower extract assisted by ultrasound method and its antibacterial activity. *Recent Patents on Nanotechnology*.
- Firdaus, N., Viquar, U., Husain, G.M., Zakir, M., 2022. Preclinical toxicity study of a *Datura metel* L. based Unani formulation *Habb-e-Zeequn Nafas* used in the treatment of Asthma. *Phytomed. Plus* 100303.
- Ghani, A., 2003. *Medicinal plants of Bangladesh with chemical constituents and uses*. Asiatic Society of Bangladesh. Nimtali, Dhaka, pp. 42–48.
- Guo, R., Liu, Y., Xu, Z.P., Xia, Y.G., Yang, B.Y., Kuang, H.X., 2018. Withanolides from the leaves of *Datura metel* L. *Phytochemistry* 155, 136–146.
- Guo, R., Liu, Y., Pan, J., Guan, W., Yang, B.Y., Kuang, H.X., 2019. A new sesquiterpenoid with cytotoxic and anti-inflammatory activity from the leaves of *Datura metel* L. *Nat. Prod. Res.* 28, 1–7.
- Han, X.L., Wang, H., Zhang, Z.H., Tan, Y., Wang, J.H., 2015. Study on Chemical Constituents in Seeds of *Datura metel* from Xinjiang. *Zhong Yao Cai* 38, 1646–1648.
- Hanif, S., Jabeen, K., Akhtar, N., Iqbal, S., 2022. GC-MS analysis & antifungal activity of *Datura metel* L. against *Rhizoctonia solani* Kuhn. *Anais da Academia Brasileira de Ciências* 94.
- Ibrahim, M., Siddique, S., Rehman, K., Husnain, M., Hussain, A., Akash, M.S.H., Azam, F., 2018. Comprehensive analysis of phytochemical constituents and

- ethnopharmacological investigation of genus *Datura*. *Crit. Rev. Eukaryot.* 28 (3), 223–283.
- Imo, C., Arowora, K.A., Ezeonu, C.S., Yakubu, O.E., Nwokwu, C.D., Azubuike, N.C., Sallah, Y.G., 2019. Effects of ethanolic extracts of leaf, seed and fruit of *Datura metel* L. on kidney function of male albino rats. *J. Tradit. Complement. Med.* 9 (4), 271–277.
- Iqbal, S., Sivaraj, C., Gunasekaran, K., 2017. Antioxidant and Anticancer Activities of Methanol Extract of Seeds of *Datura stramonium* L. *Free Radic. Antioxid.* 7, 184–189.
- Ishola, A.O., Imam, A., Ajao, M.S., 2021. Effects of datumetine on hippocampal NMDAR activity. *Toxicol. Rep.* 8, 1131–1142.
- Islam, M.T., Streck, L., Paz, M.F.C.J., Sousa, J.M.C., Alencar, M.V.O.B., Mata, A.M.O.F., Carvalho, R.M., Jose Santos, J.V.O., Silva-Junior, A.A., Ferreira, P.M.P., Melo-Cavalcante, A.A.C., 2016. Preparation of phytol-loaded nanoemulsion and screening for antioxidant capacity. *Int. Arch. Med.* 9 (70), 1–15.
- Jaafar, F.R., Ajeena, S.J., Mehdy, S.S., 2018. Anti-Inflammatory impacts and analgesic activity of aqueous extract *datura innoxia* leaves against induced pain and inflammation in mice. *J. Entomol. Zool. Stud.* 6, 1894–1899.
- Javaid, A., Shafique, S., Shafique, S., 2008. Herbicidal activity of *Datura metel* L. against *Phalaris minor* Retz. *Pak. J. Weed Sci. Res.* 14, 209–220.
- Javaid, A., Shafique, S., Shafique, S., 2010. Herbicidal activity of *Withaniasomnifera* against *Phalaris minor*. *Nat. Prod. Res.* 24, 1457–1468.
- Kam, P.C.A., Liew, S., 2002. Traditional Chinese herbal medicine and anaesthesia. *Anaesthesia* 57, 1083–1089.
- Khalili, N.R., Atyabi, S.M., 2004. Evaluation of analgesic effect of *Datura stramonium* seed extract in hot plate and formalin tested on male rats. *Iranian J. Med. Aromatic Plants* 20, 309–322.
- Ko, R.J., Ko, R., 1999. Causes, epidemiology, and clinical evaluation of suspected herbal poisoning. *J. Toxicol. Clin. Toxicol.* 37, 697–708.
- Krishna Murthy, B., Nammi, S., Kota, M.K., Krishna Rao, R.V., Koteswara Rao, N., Annapurba, A., 2004. Evaluation of hypoglycemic and antihyperglycemic effects of *Datura metel* (Linn.) seeds in normal and alloxan-induced diabetic rats. *J. Ethnopharmacol.* 91, 95–98.
- Liu, Y., Jiang, H.B., Liu, Y., Algradi, A.M., Naseem, A., Zhou, Y.Y., She, X., Yang, B.Y., Kuang, H.X., 2020. New indole alkaloids from the seeds of *Datura metel* L. *Fitoterapia* 146, 104726.
- Liu, Y., Pan, J., Sun, Y.P., Wang, X., Liu, Y., Yang, B.Y., Kuang, H.X., 2020a. Immunosuppressive withanolides from the flower of *Datura metel* L. *Fitoterapia* 141, 104468.
- Liu, Y., Tan, J.Y., Zou, H.D., Qi, Z.T., Naseem, A., Pan, J., Guan, W., Yang, B.Y., Kuang, H.X., 2021a. Seven undescribed steroids from the leaves of *Datura metel* L. *Steroids* 173, 108877.
- Liu, Y., Wu, D.D., Zhou, Y.Q., Wu, J.T., Qi, Z.T., Algradi, A.M., Pan, J., Guan, W., Yang, B. Y., Kuang, H.X., 2021b. A new ent-kaurane diterpenoid from the pericarps of *Datura metel*. *J. Asian Nat. Prod. Res.* 1–7.
- Liu, Y., Wu, D.D., Luo, Y.M., Liu, G.Z., Yang, C.L., Algradi, M., Wu, J.T., Yang, B.Y., Kuang, H.X., 2022. New sesquiterpenoid and aliphatic glycoside from the roots of *Datura metel* L. *Phytochem. Lett.* 50, 15–20.
- Maehara, S., Yamane, C., Kitamura, C., Hinokuma, M., Hata, T., 2020. High ophiobolin A production in endophytic fungus *Bipolaris* sp. associated with *Datura metel*. *Nat. Prod. Res.* 34 (20), 2990–2992.
- Mai, N.T., Cuc, N.T., Anh, H.L., Nhiem, N.X., Tai, B.H., Minh, C.V., Quang, T.H., Kim, K. W., Kim, Y.C., Oh, H., Kiem, P.V., 2017a. Steroidal saponins from *Datura metel*. *Steroids* 121, 1–9.
- Mai, N.T., Cuc, N.T., Anh, H.L.T., Nhiem, N.X., Tai, B.H., Yen, P.H., Quang, T.H., Van Minh, C., Kim, K.W., Kim, Y.C., Oh, H., 2017b. Two new guaiane sesquiterpenes from *Datura metel* L. with anti-inflammatory activity. *Phytochem. Lett.* 19, 231–236.
- Manickam, M., Sinha-Bagchi, A., Sinha, S.C., Gupta, M., Rays, A.B., 1993. Withanolides of *Datura fastuosa* leaves. *Phytochemistry* 34, 868–870.
- Manpreet, K., Anita, S., Piyush, G., 2017. Phytochemical and pharmacological study of *datura*: a review. *Int. J. Pharm. Sci. Res.* 113–118.
- Mbida, H., Tsala, D.E., Aboubakar, S., Habtemariam, S., Edmond, J.J., Bakwo, E.F., Minkande, J.Z., 2022. Antioxidant Activity of Aqueous Extract of Leaves and Seeds of *Datura metel* (Solanaceae) in Frog's Heart Failure Model. *Evid.-based Complement. Altern. Med.* 2022.
- Monira, K.M., Munan, S.M., 2012. Review on *Datura metel*: a potential medicinal plant. *Global J. Res. Med. Plants Indigenous Med.* 1, 123.
- Murch, S.J., Alan, A.R., Cao, J., Saxena, P.K., 2009. Melatonin and serotonin in flowers and fruits of *Datura metel* L. *J. Pineal Res.* 47 (3), 277–283.
- Nethradevi, C., Sivakumar, P., Renganathan, S., 2012. Green synthesis of silver nanoparticles using *Datura metel* flower extract and evaluation of their antimicrobial activity. *Int. J. Nanomat. Biostruct.* 2 (2), 16–21.
- Nkwe, D.O., Lotshwao, B., Rantong, G., Matshwele, J., Kwape, T.E., Masisi, K., Gaobotse, G., Hefferon, K., Makhzoum, A., 2021. Anticancer mechanisms of bioactive compounds from *Solanaceae*: an update. *Cancers* 13 (19), 4989.
- Ogunmoyole, T., Adeyeye, R.I., Olatilu, B.O., Akande, O.A., Agunbiade, O.J., 2019. Multiple organ toxicity of *datura stramonium* seed extracts. *Toxicol. Rep.* 2019 (6), 983–989.
- Pan, Y., Wang, X., Hu, X., 2007. Cytotoxic withanolides from the flowers of *Datura metel*. *J. Natl. Prod.* 70, 1127–1132.
- Pandiarajan, G., Govindaraj, R., Makesh, B.K., Sankarasivaraman, K., 2012. Antifertility activity in the acetone extracts of *Datura metel*, L seeds on female mouse. *J. Pharmacogenomics Pharmacoproteomics* 3, 111.
- Panter, K.E., Welch, K.D., Gardner, D.R., Green, B.T., 2013. Poisonous plants: effects on embryo and fetal development. *Birth Defects. Res. C. Embryo. Today.* 99, 223–234.
- Patachia, S., Croitoru, C., 2016. Biopolymers and biotech admixtures for eco-efficient construction materials.
- Pirintoss, S., Panagiotopoulos, A., Bariotakis, M., Daskalakis, V., Lionis, C., Sourvinos, G., Karakasiliotis, I., Kampa, M., Castanas, E., 2022. From traditional ethnopharmacology to modern natural drug discovery: a methodology discussion and specific examples. *Molecul.* 27 (13), 4060.
- Prasathkumar, M., Anisha, S., Khusro, A., Essa, M.M., Chidambaram, S.B., Qoronfleh, M. W., Sadhasivam, S., Sahibzada, M.U.K., Alghamdi, S., Almeahmadi, M., Abdulaziz, O., 2022. Anti-pathogenic, anti-diabetic, anti-inflammatory, antioxidant, and wound healing efficacy of *Datura metel* L. leaves. *Arab. J. Chem.* 15 (9), 104112.
- Preissel, U., Preissel, H.-G. (Eds.), 2002. *Brugmansia and Datura: Angel's Trumpets and Thorn Apples*. Firefly Books, Buffalo, NY, pp. 106–129. ISBN 1-55209-598-3.
- Price, D.D., 1999. Psychological mechanisms of pain and analgesia. IASP Press.
- Qin, Z., Zhang, J., Chen, L., Liu, S.X., Zhao, H.F., Mao, H.M., Zhang, H.Y., Li, D.F., 2021. Anti-inflammatory active components of the roots of *Datura metel* L. *J. Asian Nat. Prod. Res.* 23 (4), 392–398.
- Rahmatullah, M., Azam, M.N.K., Mollik, M.A.H., Hasan, M.M., Hassan, A.I., Jahan, R., Jamal, F., Nasrin, D., Ahmed, R., Rahman, M.M., Khatun, M.A., 2010. Medicinal plants used by the Kavirajes of Daulatdia Ghat, Kushtia district, Bangladesh. *Am. Eurasian J. Sustain. Agric.* 4, 219–229.
- Rajesh, Sharma, G.L., 2002. Studies on antimycotic properties of *Datura metel*. *J. Ethnopharmacol.* 80, 193–197.
- Ramadan, M.F., Zayed, R., El-Shamy, H., 2007. Screening of bioactive lipids and radical scavenging potential of some solanaceae plants. *Food Chem.* 103 (3), 885–890.
- Rauf, A., Maione, F., Uddin, G., Raza, M., Siddiqui, B.S., Muhammad, N., Shah, S.U.A., Khan, H., De Feo, V., Mascolo, N., 2016. Biological evaluation and docking analysis of daturaolone as potential cyclooxygenase inhibitor. *Evid.-based Complement. Altern. Med.* 2016.
- Rehman, M.U., Wali, A.F., Ahmad, A., Shakeel, S., Rasool, S., Ali, R., Rashid, S.M., Madkhali, H., Ganaie, M.A., Khan, R., 2019. Neuroprotective strategies for neurological disorders by natural products: an update. *Curr. Neuropharmacol.* 17 (3), 247–267.
- Rivera, R., Jacobson, I., Grimes, D., 1999. The mechanism of action of hormonal contraceptives and intrauterine contraceptive devices. *Am. J. Obstet. Gynecol.* 181 (5), 1263–1269.
- Roy, S., Pawar, S., Chowdhary, A., 2016. Evaluation of in vitro cytotoxic and antioxidant activity of *datura metel* Linn. and *cynodon dactylon* Linn. extracts. *Pharmacogn. Res.* 8, 123.
- Samuelsson, B., 1987. An elucidation of the arachidonic acid cascade: discovery of prostaglandins, thromboxane and leukotrienes. *Drugs* 33 (Suppl 1), 2–9.
- Sangeetha, S., Deepa, M., Sugitha, N., Mythili, S., Sathivelu, A., 2014. Antioxidant activity and phytochemical analysis of *Datura metel*. *Int. J. Drug Dev. Res.* 6 (4), 46–53.
- Schultes, R.E., Hofmann, A., 1979. *Plants of the gods: origins of hallucinogenic use*. 149.
- Shuping, D.S.S., Eloff, J.N., 2017. The use of plants to protect plants and food against fungal pathogens: a review. *Afric. J. Tradit. Complement. Altern. Med.* 14, 120–127.
- Siddiqui, S., Sultana, N., Ahmad, S.S., Haider, S.I., 1987. A novel withanolide from *Datura metel*. *Phytochemistry* 26, 2641–2643.
- Singh, V., Singh, R., 2008. Effect of *Datura metel* seed methanol extract and its fractions on the biology and ovipositional behaviour of *Helicoverpa armigera*. *J. Med. Aromatic Plant Sci.* 30, 157–163.
- Su, Y., Zhang, F., Wu, L., Kuang, H., Wang, Q., Cheng, G., 2022. Total withanolides ameliorates imiquimod-induced psoriasis-like skin inflammation. *J. Ethnopharmacol.* 285, 114895.
- Tan, J.Y., Liu, Y., Cheng, Y.G., Sun, Y.P., Liu, Y., Huang, J., Guo, S., Liu, G.Z., Kuang, H. X., Yang, B.Y., 2020a. Daturmetesides AE, five new ergostane-type C28 sterols from the leaves of *Datura metel* L. *Steroids* 156, 108583.
- Tan, J.Y., Liu, Y., Cheng, Y.G., Sun, Y.P., Pan, J., Yang, S.H., Kuang, H.X., Yang, B.Y., 2020b. Anti-inflammatory sesquiterpenoids from the leaves of *Datura metel* L. *Fitoterapia* 142, 104531.
- Tan, J.Y., Liu, Y., Cheng, Y.G., Sun, Y.P., Li, X.M., Guan, W., Pan, J., Yang, B.Y., Kuang, H.X., 2021. Seven new glycosides from the leaves of *Datura metel* L. *Nat. Prod. Res.* 36 (1), 295–304.
- Tariq, A., Mussarat, S., Adnan, M., 2015. Review on ethnomedicinal, phytochemical and pharmacological evidence of Himalayan anticancer plants. *J. Ethnopharmacol.* 164, 96–119.
- Tripathi, Y.C., Prabhu, V.V., Pal, R.S., Mishra, R.N., 1996. Medicinal plants of Rajasthan in Indian system of medicine. *Ancient Sci. Life* 15, 190.
- Varma, A., Padh, H., Shrivastava, N., 2011. Andrographolide: a new plant-derived antineoplastic entity on horizon. *Evid.-based Complement. Altern. Med.* p. 2011.
- Vimal, A.S., Suseela, L., 2010. In vitro antioxidant activity and wound healing activity of the alcoholic extract of the alcoholic extract of the ariel parts of *Datura fastuosa* Linn. *J. Pharm. Res.* 2, 1176–1179.
- Wang, X.Y., He, J., Bai, H.J., Gao, P.Y., Gan, H., Yang, T., Zhang, W.K., Xu, J.K., 2019. Daturanolide A-C, Three New Withanolides from *Datura metel* L. and Their Cytotoxic Activities. *Chem. Biodivers.* 16, e1900004.
- Wang, S.Y., Liu, Y., Li, X.M., Algradi, A.M., Jiang, H., Sun, Y.P., Guan, W., Pan, J., Kuang, H.X., Yang, B.Y., 2021. Discovery of Active Ingredients Targeted TREM2 by SPR Biosensor-UPLC/MS Recognition System, and Investigating the Mechanism of Anti-Neuroinflammatory Activity on the Lignin-Amides from *Datura metel* Seeds. *Molecules* 26 (19), 5946.
- Wang, Q.H., Xiao, H.B., Yang, B.Y., Yao, F.Y., Kuang, H.X., 2008. Studies on pharmacological actions of the effective parts for psoriasis in *Flos Daturae* (I). *Chinese J. Exp. Trad. Med. Formulae* 14, 48–51.
- Wannang, N.N., Ndukwe, H.C., Nnabuife, C., 2009. Evaluation of the analgesic properties of the *Datura metel* seeds aqueous extract. *J. Med. Plants Res.* 3, 192–195.

- Wei, Z., Li, T., Sun, Y., Su, H., Zeng, Y., Wang, Q., Kuang, H., 2021. Daturaturin A, a withanolide in *Datura metel* L., induces HaCaT autophagy through the PI3K-Akt-mTOR signaling pathway. *Phytother. Res.* 35 (3), 1546–1558.
- Wijesekara, K.B., Iqbal, M., 2021. Induction of Haploid Embryos in *Datura metel* by Anther Culture. In: *Doubled Haploid Technology*. Humana, New York, NY, pp. 327–336.
- Wild, D., von Maltzahn, R., Brohan, E., Christensen, T., Clauson, P., Gonder-Frederick, L., 2007. A critical review of the literature on fear of hypoglycemia in diabetes: implications for diabetes management and patient education. *Patient Educ. Counsel.* 68 (1), 10–15.
- Xia, C., Liu, Y., Qi, H., Niu, L., Zhu, Y., Lu, W., Xu, X., Su, Y., Yang, B., Wang, Q., 2019. Characterization of the metabolic fate of *Datura metel* seed extract and its main constituents in rats. *Front. Pharmacol.* 10, 571.
- Xiang, Y., Guo, Z., Zhu, P., Chen, J., Huang, Y., 2019. Traditional Chinese medicine as a cancer treatment: Modern perspectives of ancient but advanced science. *Cancer Med.* 8 (5), 1958–1975.
- Yamazaki, Z., Tagaya, I., 1980. Antiviral effects of atropine and caffeine. *J. General Virol.* 50, 429–431.
- Yang, B., Wang, Q., Xia, Y., Feng, W., Kuang, H., 2007. Withanolide Compounds from the Flower of *Datura metel* L. *Helvetica Chim. Acta* 90 (8), 1522–1528.
- Yang, B.Y., Xia, Y.G., Wang, Q.H., Dou, D.Q., Kuang, H.X., 2010a. Baimantuoluosides D-G, four new withanolide glucosides from the flower of *Datura metel* L. *Arch. Pharm. Res.* 33 (8), 1143–1148.
- Yang, B.Y., Xia, Y.G., Wang, Q.H., Dou, D.Q., Kuang, H.X., 2010b. Two new amide alkaloids from the flower of *Datura metel* L. *Fitoterapia* 81 (8), 1003–1005.
- Yang, B.Y., Zhou, Y.Q., Liu, Y., Lu, Z.K., Kuang, H.X., 2017. Withanolides as potential immunosuppressive agents against RAW264.7 cells from the pericarps of *Datura metel*. *Nat. Prod. Commun.* 12, 1934578X1701200705.
- Yang, B.Y., Luo, Y.M., Liu, Y., Yin, X., Zhou, Y.Y., Kuang, H.X., 2018. New lignans from the roots of *Datura metel* L. *Phytochem. Lett.* 28, 8–12.
- Yang, B.Y., Jiang, H.B., Liu, Y., Chen, J., Kuang, H.X., 2020. Steroids from the seeds of *Datura metel*. *J. Asian Nat. Prod. Res.* 22 (3), 257–263.
- Yu, M.Y., Zhao, G.T., Liu, J.Q., Khan, A., Peng, X.R., Zhou, L., Dong, J.R., Li, H.Z., Qiu, M. H., 2017. Withanolides from aerial parts of *Nicandraphysalodes*. *Phytochemistry* 137, 148–155.
- Yulizar, Y., Kusriani, E., Apriandanu, D.O.B., Nurdini, N., 2020. *Datura metel* L. Leaves extract mediated CeO₂ nanoparticles: synthesis, characterizations, and degradation activity of DPPH radical. *Surf. Interfaces* 19, 100437.
- Yusuf, M., Chowdhury, J.U., Wahab, M.A., Begum, J., 1994. Medicinal plants of Bangladesh. Bangladesh Council of Scientific and Industrial Research, Dhaka, Bangladesh, p. 192.
- Yusuf, M., Begum, J., Hoque, M.N., Chowdhury, 2009. Medicinal plants of Bangladesh. BCSIR Chittagong, Bangladesh, p. 794.